


COLOMBIA

Leishmaniasis cutaneous and mucosal 2015



LCIc



Incidence
(100.000 pop.) **33.6**

Transmission
Leishmaniasis
Composite Indicator
(LCIc) **Intense**

Occurrence

7541
NEW CASES

7464 cutaneous (99%)
77 mucosal (1%)



AGE GROUP

4.8% < 5 years old
6.1% ≥ 5 < 10 years old
88.8% > 10 years old
0.3% Unknown

10.9% of children
under 10 years old



SEX

76.4% Male
23.6% Female

57% of men over
20 years old



CROSS-BORDER

Shares borders with
5 countries in **42**
municipalities

6.8% of the cases
at the borders



SURVEILLANCE AND CONTROL

46% of the population
in transmission areas

Individual Mandatory
Notification

Vector chemical control



LEISHMANIA SPECIES

L. amazonensis
L. braziliensis
L. mexicana
L. colombiensis
L. guyanensis
L. panamensis



LUTZOMYIA VECTORS

L. flaviscutellata, *L. colombiana*, *L. spinicrassa*, *L. pia*, *L. towsendi*, *L. hartmanni*, *L. umbratilis*, *L. longiflocosa*, *L. trapedoi*, *L. panamensis*, *L. yulli yulli*, *L. cruciata*, *L. columbiana*, *L. gomezi*



PATIENT ASSISTANCE

Free treatment at
healthcare facilities

Laboratory confirmation:
100%

Cure Rate: **No
Information**



AVAILABLE MEDICINES

Meglumine
antimoniate

Source:

SisLeish - PAHO/WHO - Data provided by the Ministries of Health -
National Leishmaniasis Programs of each country
Access on: December, 2016